

PIPRA AUREOLA.



CHARACTER GENERICUS.

Rostrum capite brevius, basi subtrigonum, integerrimum, apice incurvum.

Pedes gressorii.

Lin. Syst. Nat. p. 338.

CHARACTER SPECIFICUS, &c.

PIPRA NIGRA, capite pectoreque coccineis aut luteis.

PIPRA NIGRA, capite pectoreque coccineis, remigibus antrorsum macula alba.

Lin. Syst. Nat. p. 339.

PARUS AUREOLA.

Lin, Syst. Nat. ed. 10. p. 191.

PARUS RUBER et NIGER.

Edke. av. 2. p. 109. t. 261.

MANACUS RUBER.

Brieff. av. 4. p. 452. t. 34. f. 3.

Coloribus variat *Pipra aureola*; interdum enim capite pectoreque coccineis, interdum luteis conspicitur. Americam incolit calidiorem et in Cayana præcipue invenitur. Magnitudinem veram exprimit tabula.



A. 1.

London, Published July 1st 1796 by F. & J. Wadsworth & Co. No. 55. Brewer Street.

THE
RED-AND-BLACK MANAKIN.



GENERIC CHARACTER.

Bill shorter than the head, somewhat triangular at the base, bent at tip.

Feet gressorial.

SPECIFIC CHARACTER.

BLACK MANAKIN with the head and breast either scarlet or gold-yellow.

The RED and BLACK MANAKIN.

Edw. t. 261. f. 2.

Le MANAKIN rouge.

Buff. ois. 4. p. 415.

Le MANAKIN orangé.

Buf. ois. 4. p. 417.

Pl. enl. 34. f. 3. & 302. f. 2.

The *Pipra aurcola* varies in color, being sometimes seen with the head and breast of a rich scarlet, at other times of a gold-yellow. It is an inhabitant of the hotter parts of America, and is principally found in Cayenne. The plate represents it in its natural size.